

EGOROV, LUSIN and HAUSSDORFF.

Topic 1. Lusin Thm.

Let μ be a Borel regular measure in $\mathbb{R}^n$. $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be μ -measurable. A is μ -measurable. $\mu(A) < +\infty$.

Fix $\varepsilon > 0$. Then there exists a compact set $K \subseteq A$ such that.

(i) $\mu(A \setminus K) < \varepsilon$ (ii) $f|_K$ is cts.

Proof. Step 1. Construct a compact subset of A .

Fix $i \in \mathbb{N}$. Let $B_{ij} \subseteq \mathbb{R}^m$ be disjoint subsets s.t. $\bigcup_{j=1}^{\infty} B_{ij} = \mathbb{R}^m$. $\text{diam } B_{ij} < \frac{1}{i}$

Let $A_{ij} = f^{-1}(B_{ij}) \cap A$. Then A_{ij} is μ -measurable. $A = \bigcup_{j=1}^{\infty} A_{ij}$

Write $\nu = \mu|_A$ (μ restrict to A) i.e. $\nu(x) = \mu(x \cap A)$ ν is Radon measure (finite, Borel regular).

There exists $K_{ij} \subseteq A_{ij}$ compact. $\nu(A_{ij} \setminus K_{ij}) < \frac{\varepsilon}{2^{i+j}}$

$$\mu(A \setminus \bigcup_{j=1}^{\infty} K_{ij}) = \nu(A \setminus \bigcup_{j=1}^{\infty} K_{ij}) = \nu(\bigcup_{j=1}^{\infty} A_{ij} \setminus \bigcup_{j=1}^{\infty} K_{ij}) \leq \nu(\bigcup_{j=1}^{\infty} (A_{ij} \setminus K_{ij})) \leq \sum_{j=1}^{\infty} \frac{\varepsilon}{2^{i+j}} = \frac{\varepsilon}{2^i}$$

$$\bigcup_{j=1}^{\infty} A_{ij} \setminus \bigcup_{j=1}^{\infty} K_{ij} \subseteq \bigcup_{j=1}^{\infty} (A_{ij} \setminus K_{ij})$$

Warning: $\bigcup_{j=1}^{\infty} K_{ij}$ may not be compact. Since $\lim_{N \rightarrow \infty} \mu(A \setminus \bigcup_{j=1}^N K_{ij}) = \mu(A \setminus \bigcup_{j=1}^{\infty} K_{ij}) < \frac{\varepsilon}{2^i}$. There exists $N_i \in \mathbb{N}$ s.t.


$$\mu(A \setminus \bigcup_{j=1}^{N_i} K_{ij}) < \frac{\varepsilon}{2^{i+1}} \quad \text{Now finite union of compact set is compact. Call } D_i := \bigcup_{j=1}^{N_i} K_{ij}, \text{ compact.}$$

Step 2. Construct a cts. function.

Fix $b_{ij} \in B_{ij} \subseteq \mathbb{R}^m$. $\text{diam}(B_{ij}) < \frac{1}{i}$. Define $g_i: D_i \rightarrow \mathbb{R}^m$. $g_i(x) = b_{ij}$ for $x \in K_{ij}$, $1 \leq j \leq N_i$

$$= \sum_{j=1}^{N_i} b_{ij} \mathbf{1}_{K_{ij}}$$

$|f(x) - g_i(x)| < \frac{1}{i}$ since if $x \in D_i$, $f(x) \in B_{ij}$ for some j .



Let $K := \bigcap_{i=1}^{\infty} D_i$ still compact. $\mu(A \setminus K) = \mu(A \setminus \bigcap_{i=1}^{\infty} D_i) \leq \sum_{i=1}^{\infty} \mu(A \setminus D_i) = \sum_{i=1}^{\infty} \frac{\varepsilon}{2^{i+1}} < 2\varepsilon$. (not tight)

$g_i \Rightarrow f$ uniformly on K . $g := f|_K$ is cts.

Topic 2. Egorov Thm.

$f_k: X \rightarrow \mathbb{R}^n$ be μ -measurable functions. $f_k \rightarrow f$ a.e. on X . $\mu(X) < +\infty$. Then there exists $B \subseteq A$ s.t.

(i) $\mu(A \setminus B) < \varepsilon$. (ii) $f_k \rightarrow f$ on B .

$$\mu\left(\bigcap_{i=1}^{\infty} \bigcup_{j=1}^{\infty} \bigcap_{k \geq j} \left\{x \in X : |f_k(x) - f(x)| < \frac{1}{i}\right\}\right) = \mu(X).$$

Annotations: $\forall \varepsilon > 0, \exists N, \forall n \geq N, \exists \delta > 0$. $|f(x) - f_n(x)| < \varepsilon$.

Proof. $C_{i,j} = \bigcup_{k=j}^{\infty} \left\{x \in X : |f_k(x) - f(x)| > \frac{1}{2^i}\right\}$, i, j fixed.

So $C_{i,j+1} \subseteq C_{i,j}$ decreasing set. So $\lim_{j \rightarrow \infty} \mu(C_{i,j}) = \mu\left(\bigcap_{j=1}^{\infty} C_{i,j}\right) = 0$ by monotone conv.

$\Rightarrow \exists N(i) \in \mathbb{N}$ s.t. $\mu(C_{i, N(i)}) < \frac{\varepsilon}{2^i}$. Let $B^c = \bigcup_{i=1}^{\infty} C_{i, N(i)}$. $\mu(B^c) \leq \sum_{i=1}^{\infty} \frac{\varepsilon}{2^i} = \varepsilon$.

$\forall x \in B$: $x \in \bigcap_{i=1}^{\infty} C_{i, N(i)}^c$, so $|f(x) - f_n(x)| < \frac{1}{2^i}$ if $n > N(i)$. $\Leftrightarrow$ Unif. conv.

Alternative similar proof: $f_n \xrightarrow{a.e.} f \Leftrightarrow \mu\left[\left(\bigcap_{k=1}^{\infty} \bigcup_{n \geq k} \bigcap_{m \geq n} \left\{x : |f_m(x) - f_n(x)| < \frac{1}{k}\right\}\right)^c\right] = 0$

$\Leftrightarrow \mu\left(\bigcup_{k=1}^{\infty} \bigcap_{n \geq 1} \bigcup_{m \geq n} \left\{|f_m(x) - f_n(x)| \geq \frac{1}{k}\right\}\right) = 0.$

$\Rightarrow \mu\left(\bigcup_{n \geq 1} \bigcap_{m \geq n} \left\{|f_m(x) - f_n(x)| \geq \frac{1}{k}\right\}\right) = 0 \quad \forall k.$

$\lim_{N \rightarrow \infty} \mu\left(\bigcup_{n \geq N} \left\{|f_n(x) - f(x)| \geq \frac{1}{k}\right\}\right) = 0$ *EM decreasing set sequence* $\mu(X) < +\infty$.

$\Leftrightarrow \exists N_k, \mu\left(\bigcup_{n \geq N_k} \left\{|f_n(x) - f(x)| \geq \frac{1}{k}\right\}\right) < \frac{\varepsilon}{2^{k+1}} \quad \mu(E_{N_k}) < \frac{\varepsilon}{2^{k+1}} \quad E := \bigcup_{k=1}^{\infty} E_{N_k} \Rightarrow \mu(E) \leq \sum_{k=1}^{\infty} \mu(E_{N_k}) = \frac{\varepsilon}{2} < \varepsilon$

Claim: $f_n \rightarrow f$ on $X \setminus E$.

Topic 3 Hausdorff measure.

(i) Let $A \subseteq \mathbb{R}^n$. $0 \leq s < +\infty$ $0 < \delta \leq +\infty$.

$H_s^\delta(A) = \inf \left\{ \sum_{j=1}^{\infty} \alpha(s) \left(\frac{\text{diam } C_j}{2}\right)^s \mid A \subseteq \bigcup_{j=1}^{\infty} C_j, \text{diam } C_j < \delta \right\}$ Here $\alpha(s) = \frac{\pi^{\frac{s}{2}}}{\Gamma(\frac{s}{2} + 1)}$

(ii) The Hausdorff measure of A

$H^s(A) := \sup_{\delta > 0} H_s^\delta(A) = \lim_{\delta \rightarrow 0} H_s^\delta(A).$

R.M.K. 1) $\Gamma(s) = \int_0^\infty e^{-x} x^{s-1} dx$ ($0 < s < \infty$) Gamma function.

2) $\text{vol}(B(x, r)) = \alpha(n) r^n$. Where $B(x, r)$ is the ball in $\mathbb{R}^n$ with radius r .

3) $H_s^\delta \downarrow$ in δ



Theorem. Hausdorff measure H^s is Borel regular. for $0 \leq s < +\infty$

Proof. (i) Borel 集都可测. See Evans. 304 → Homework 2

(ii) Borel regularity: Given A . $\exists$ Borel set $B \subseteq A$ $H^s(B) = H^s(A)$.

$$H^s_\delta(A) = \inf \left\{ \sum_{j=1}^{\infty} \alpha(s) \left(\frac{\text{diam } C_j}{2} \right)^s \mid A \subseteq \bigcup_{j=1}^{\infty} C_j, \text{diam } C_j < \delta \right\}$$
$$= \inf \left\{ \sum_{j=1}^{\infty} \alpha(s) \left(\frac{\text{diam } \bar{C}_j}{2} \right)^s \mid A \subseteq \bigcup_{j=1}^{\infty} \bar{C}_j, \text{diam } \bar{C}_j < \delta \right\} \quad \bar{C}_j \text{ is the closure of } C_j.$$

$\Rightarrow$ We could require C_j be closed.

$$H^s_\delta(A) = \inf \left\{ \sum_{j=1}^{\infty} \alpha(s) \left(\frac{\text{diam } C_j}{2} \right)^s \mid A \subseteq \bigcup_{j=1}^{\infty} C_j, \text{diam } C_j < \delta, C_j \text{ closed} \right\}$$

Let $A \subseteq \mathbb{R}^n$. $H^s(A) < +\infty \Rightarrow H^s_\delta(A) < +\infty$.

Choose closed set $\{C_{kj}\}_{j=1}^{\infty}$ $\text{diam } C_{kj} < \frac{1}{k}$ covering A . and

$$\sum_{j=1}^{\infty} \alpha(s) \left(\frac{\text{diam } C_{kj}}{2} \right)^s < \underbrace{H^s_{\frac{1}{k}}(A)}_{\uparrow \delta} + \underbrace{\frac{1}{k}}_{\uparrow \varepsilon} = \sum \text{diam} \dots H^s_\delta + \varepsilon.$$

Let $A_k = \bigcup_{j=1}^{\infty} C_{kj}$ $B = \bigcap_{k=1}^{\infty} A_k$ Claim: $A \subseteq B$. Borel $H^s(A) = H^s(B)$
trivial

Since $B = \bigcap_{k=1}^{\infty} A_k$, $B \subseteq A_k = \bigcup_{j=1}^{\infty} C_{kj}$ for fix k .

$$\Rightarrow H^s_{\frac{1}{k}}(B) \leq \sum_{j=1}^{\infty} \alpha(s) \left(\frac{\text{diam}(C_{kj})}{2} \right)^s \leq H^s_{\frac{1}{k}}(A) + \frac{1}{k}.$$

let $k \rightarrow +\infty$ $H^s(B) \leq H^s(A)$ } equality holds

Since $A \subseteq A_k \forall k, \Rightarrow A \subseteq B$. $H^s(B) \geq H^s(A)$

Thm. properties of H^s .

(i) $H^0 \equiv$ counting measure on $\mathbb{R}^n$.

(ii) $H^s \equiv 0$ if $s > n$.

(iii) $H^1 = \lambda^1$ 1D Lebesgue on $\mathbb{R}$. ($H^n = \lambda^n$ on $\mathbb{R}^n$. but the proof is not easy)

(iv) $H^s(\alpha A) = \alpha^s H^s(A)$ for $\alpha > 0$. $A \subseteq \mathbb{R}^n$

(v) $H^s(L(A)) = H^s(A)$ where $A \subseteq \mathbb{R}^n$, $L: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is an affine isometry.

Proof: Evans. Chapter 2.

Other details: See Evans.