

Take $\{u_{nn}\}_{n=1}^{\infty}$. u_{nn} converges in $L^q(V)$

RMK. $p^* > p$, $p^* \rightarrow \infty$ as $p \rightarrow n$.

$$\Rightarrow W^{1,p}(U) \hookrightarrow L^{p^*}(U) \quad \forall p \in [1, \infty).$$

For $n < p < \infty$

Poincaré Inequality

$$(u)_U = \int_U u dy = \frac{1}{|U|} \int_U u dy$$

Thm. U bounded, open, connected subset of $\mathbb{R}^n$, with C^1 boundary. $1 \leq p \leq \infty$.

$$\|u - (u)_U\|_{L^p(U)} \lesssim_{n,p,U} \|Du\|_{L^p(U)} \quad \forall u \in W^{1,p}$$

RMK if $|U| < \infty$, then can control u in terms of its gradient.

Proof. We argue by contradiction. Suppose the ineq. is false. Fix $U \subseteq \mathbb{R}^n$. For each $k=1,2,\dots$ we can find

$$u_k \in W^{1,p}(U) \text{ s.t. } \|u_k - (u_k)_U\|_{L^p} > k \|Du_k\|_{L^p}$$

$$\text{Re-normalize } v_k = \frac{u_k - (u_k)_U}{\|u_k - (u_k)_U\|_{L^p(U)}}, \quad k=1,2,\dots$$

$$\Rightarrow \begin{cases} \|v_k\|_{L^p} = 1 \\ (v_k)_U = 0 \\ \|Dv_k\|_{L^p} \leq \frac{1}{k} \end{cases} \Rightarrow v_k \text{ bounded in } W^{1,p}(U)$$

By using Rellich-Kondrachov (and the remarks after)

$\exists$ a subsequence $u_{k_j} \in W^{1,p}(U)$ converging in $L^p(U)$ to $v \in L^p(U)$.

$$\begin{cases} \|u_{k_j}\|_{L^p} = 1 \\ (v_{k_j})_U = 0 \end{cases} \Rightarrow \begin{cases} \|v\|_{L^p} = 1 \\ (v)_U = 0 \end{cases}$$

$$\|Dv_{k_j}\|_{L^p} \leq \frac{1}{k_j} \xrightarrow{j \rightarrow \infty} 0 \implies Dv = 0$$

Proof of the " $\Rightarrow$ ": Take $\phi \in C_c^\infty(U)$. By def of weak derivative:

$$-\int_U v \phi_{x_i} = -\lim_{j \rightarrow \infty} \int_U v_{k_j} \phi_{x_i} = \lim_{j \rightarrow \infty} \int_U (v_{k_j})_{x_i} \phi \leq \lim_{j \rightarrow \infty} \frac{1}{k_j} = 0$$

$$\Rightarrow Dv = 0$$

Now $v \in W^{1,p}$, $Dv = 0$ a.e. $\xrightarrow{\triangle} U \text{ open connected} \Rightarrow v \equiv \text{Const. a.e.}$

$$\text{But } \|v\|_{L^p} = 1, \quad (v)_U = 0$$

$$\downarrow \quad \downarrow \\ v \text{ cannot be } 0 \text{ a.e.} \quad v = \text{Const.} = 0$$

Contradiction.

Thm. Poincaré Ineq. for a Ball

$$\|u - (u)_{B(x,r)}\|_{L^p(B(x,r))} \lesssim r \cdot \|Du\|_{L^p(B(x,r))} \quad \forall u \in W^{1,p}(B(x,r))$$

Proof. Case $U = B(0,1)$ follows from the previous thm.

General case: Rescale $v(y) = u(x + ry)$ $y \in B(0,1)$

$$\text{Then } v \in W^{1,p}(B(0,1)) \quad \|v - (v)_{B(0,1)}\|_{L^p(B(0,1))} \lesssim \|Dv\|_{L^p(B(0,1))}$$

Now change of variable. RHS will yield a "r" factor.

RMK. Connection of BMO and $W^{1,n}$

Assume $u \in W^{1,n}(\mathbb{R}^n) \cap L^1(\mathbb{R}^n)$ ($n \geq 2$) Take $p=1$ in the previous theorem.

$$\int_{B(x,r)} |u - (u)_{B(x,r)}| \lesssim r \int_{B(x,r)} |Du|$$

$$\stackrel{\text{Hölder}}{\lesssim} r \left(\int_{B(x,r)} |Du|^n \right)^{\frac{1}{n}}$$

$$= \frac{1}{(\alpha(n))^{\frac{1}{n}}} \left(\int |Du|^n \right)^{\frac{1}{n}}$$

$$\Rightarrow u \in \boxed{\text{BMO}(\mathbb{R}^n)}$$

$$\text{Def. } [u]_{\text{BMO}(\mathbb{R}^n)} := \sup_{B(x,r) \subseteq \mathbb{R}^n} \left\{ \int_{B(x,r)} |u - (u)_{B(x,r)}| \right\}$$

Difference quotients. 差商

Assume $u: \Omega \rightarrow \mathbb{R}^n \in L^1_{loc}(\Omega)$. $V \subset\subset \Omega$.

Def. (i) The i -th difference quotient of size h

$$D_i^h u(x) := \frac{u(x + h e_i) - u(x)}{h} \quad (i=1,2,\dots,n) \quad \forall x \in V, \quad h \in \mathbb{R}, \quad 0 < |h| < \text{dist}(V, \partial\Omega)$$

$$(ii) D^h u := (D_1^h u, \dots, D_n^h u)$$

Theorem

(i) $1 \leq p < \infty$. $u \in W^{1,p}(\Omega)$ For each $V \subset\subset \Omega$, we have $\|D^h u\|_{L^p(V)} \lesssim_{\Omega, V, p, n} \|Du\|_{L^p(\Omega)}$

for each $0 < |h| < \frac{1}{2} \text{dist}(V, \partial\Omega)$

Proof of (i): Assume $u \in C^\infty(\Omega)$ then use density.

$$u(x + h e_i) - u(x) = h \int_0^1 u_{x_i}(x + t h e_i) dt$$

$$|u(x + h e_i) - u(x)| \leq |h| \int_0^1 |Du(x + t h e_i)| dt$$

$$\begin{aligned} \Rightarrow \int_V |D^h u|^p dx &\lesssim \sum_{i=1}^n \int_V \int_0^1 |Du(x + t h e_i)|^p dt dx \\ &= \sum_{i=1}^n \int_0^1 \int_V |Du(x + t h e_i)|^p dx dt \\ &\lesssim \int_\Omega |Du|^p dx \int_0^1 dt \\ &= \|Du\|_{L^p(\Omega)}^p. \end{aligned}$$

(ii) Assume $1 < p < \infty$ $u \in L^p(\Omega) \exists \text{ const. } C$ s.t.

$$\|D^h u\|_{L^p} \leq C, \quad \forall 0 < |h| < \frac{1}{2} \text{dist}(V, \partial\Omega)$$

Then: $u \in W^{1,p}(\Omega)$. $\|Du\|_{L^p(\Omega)} \leq C$.

Proof. Suppose $\|D^h u\|_{L^p(\Omega)} \leq C \quad \forall 0 < |h| < \frac{1}{2} \text{dist}(V, \partial\Omega)$. Choose $\phi \in C_c^\infty(\Omega)$ then for small h

$$\int_\Omega u(x) \left[\frac{\phi(x + h e_i) - \phi(x)}{h} \right] dx = - \int_\Omega \frac{u(x) - u(x - h e_i)}{h} \phi(x) dx$$

$$\int_\Omega u (D_i^h \phi) = - \int_\Omega (D_i^{-h} u) \phi$$

Integration by parts formula.

$\sup_h \|D_i^{-h} u\|_{L^p(\Omega)} < \infty$ Behave like Balzano - Wienerstrauss

Now since $1 < p < \infty$, there exists a function $v_i \in L^p(\mathcal{V})$ and $h_k \rightarrow 0$ s.t.

$$D_i^{h_k} u \longrightarrow v_i \text{ weakly in } L^p(\mathcal{V})$$

$$\int_{\mathcal{V}} u \phi_{x_i} = \int_{\mathcal{V}} u \phi_{x_i} = \lim_{h_k \rightarrow 0} \int_{\mathcal{V}} u D_i^{h_k} \phi$$

$$\stackrel{\substack{\text{integration} \\ \text{by parts}}}{=} - \lim_{h_k \rightarrow 0} \int_{\mathcal{V}} (D_i^{h_k} u) \phi$$

$$\stackrel{\substack{\text{weak} \\ \text{conv.}}}{=} - \int_{\mathcal{V}} v_i \phi = - \int_{\mathcal{V}} u_i \phi$$

$\Rightarrow v_i = u_{x_i}$ in the weak sense

$$\Rightarrow D u \in L^p(\mathcal{V}) \Rightarrow v \in W^{1,p}(\mathcal{V})$$

$$\left\{ \begin{array}{l} u \in L^p(\mathcal{V}) \end{array} \right.$$

* Theorem. (Characterization of $W^{1,\infty}$)

Let $\mathcal{U} \subseteq \mathbb{R}^n$ be open bounded with C^1 boundary. Then

$u: \mathcal{U} \rightarrow \mathbb{R}$ is Lipschitz continuous $\Leftrightarrow u \in W^{1,\infty}(\mathcal{U})$

Proof. First assume $\mathcal{U} = \mathbb{R}^n$, and u is compact supported.

Suppose $u \in W^{1,\infty}(\mathbb{R}^n)$. $\|Du\|_{\infty}, \|u\|_{\infty} < \infty$. Take $u^\varepsilon = \eta_\varepsilon * u$ the mollification of u . Then $u^\varepsilon \in C_c^\infty(\mathbb{R}^n)$

$u^\varepsilon \rightarrow u$ unif. as $\varepsilon \rightarrow 0$ on finite domain. $u \in W^{1,p} (p \geq n)$. Use Morrey. u is cts

Take $x, y \in \mathbb{R}^n$. $x \neq y$. $u^\varepsilon(x) - u^\varepsilon(y) = \int_0^1 \frac{d}{dt} u^\varepsilon(tx + (1-t)y) dt$ (Fund. Thm. of Calc.)

$$= \int_0^1 (Du^\varepsilon)(tx + (1-t)y) dt \cdot (x-y)$$

Does not hold for original function u .

$$\Rightarrow |u^\varepsilon(x) - u^\varepsilon(y)| \leq \|Du^\varepsilon\|_{\infty} |x-y|$$

$$\leq \|Du\|_{\infty} |x-y|.$$

Now send $\varepsilon \rightarrow 0$. use unif. convergence. we are done.

② Suppose u is Lipschitz continuous: use Diff. quotient.

$$u \text{ Lipschitz} \Rightarrow \|D_i^{-h} u\|_{\infty} \leq \text{Lipschitz const. of } u := \text{Lip}(u)$$

there exists a function $v_i \in L^{\infty}(\mathbb{R}^n)$ and a subsequence $h_k \rightarrow 0$ s.t. $D_i^{-h_k} u \xrightarrow{\text{weak}} v_i$ in $L^{\infty}_{loc}(\mathbb{R}^n)$

$$\begin{aligned} \Rightarrow \int_{\mathbb{R}^n} \phi_{x_i} \cdot u &= \lim_{h_k \rightarrow 0} \int_{\mathbb{R}^n} u \cdot D_i^{h_k} \phi = \lim_{h_k \rightarrow 0} - \int_{\mathbb{R}^n} (D_i^{-h_k} u) \phi \\ &= - \int_{\mathbb{R}^n} v_i \phi \end{aligned}$$

$\Rightarrow v_i = u_{x_i}$ in the weak sense, $i = 1, 2, \dots, n$.

$\Rightarrow u \in W^{1,\infty}(\mathbb{R}^n)$

RMK. the trick of Diff. quotient.

Classical: $D_i^h u := \frac{u(x+he_i) - u(x)}{h} \rightarrow D_i u$

Sobolev: $D_i^h u \xrightarrow{\text{weakly}} D_i u$ weak derivative

2. General Case: U bounded, ∂U is C^1 .

We first extend u to $\bar{u} := Eu$ on $\mathbb{R}^n$. Repeat the same argument. []

RMK. For any open set U , also prove: u is locally Lipschitz $\iff u \in W^{1,\infty}_{loc}(U)$

If $n < p < \infty$: $u \in W^{1,p} \xrightarrow{\text{Morrey}} u \in C^{0,1-\frac{n}{p}}$

Differentiability a.e.

Def. A function $u: \mathcal{U} \rightarrow \mathbb{R}$ is differentiable at $x_0 \in \mathcal{U}$ if $\exists a \in \mathbb{R}^n$ s.t.

$$u(y) = u(x) + a \cdot (y-x) + o(|y-x|)$$

$$\text{i.e. } \lim_{y \rightarrow x} \frac{|u(y) - u(x) - a \cdot (y-x)|}{|y-x|} = 0.$$

RMK. a is unique. Denote $Du := a$.

Theorem. (Differentiability a.e.)

Assume $u \in W_{loc}^{1,p}(\mathcal{U})$ for some $n < p \leq \infty$. Then u is differentiable a.e. in $\mathcal{U}$.

It's gradient equals its weak gradient a.e.

RMK. We identify u as its continuous version, as before.

Proof Recall Morrey for $n < p < \infty$: $|u(y) - u(x)| \lesssim r^{1-\frac{n}{p}} \left(\int_{B(x,r)} |Du(z)|^p dz \right)^{\frac{1}{p}}, y \in B(x,r)$

valid for $v \in C^1$. (By approximation, it's also true for $u \in W^{1,p}$).

If one has $\int_{B(x,r)} |Du(z)|^p dz$, then

$$|u(x) - u(y)| \lesssim r^{1-\frac{n}{p}} |Du(x)| r^{\frac{n}{p}}$$

$$\lesssim |Du(x)| r, \text{ as needed.}$$

Hence Hope: $\int_{B(x,2r)} |Du(x) - Du(z)|^p dz$ small.

Formal proof: Choose $u \in W_{loc}^{1,p}(\mathcal{U})$. Then for Lebesgue pts $x \in \mathcal{U}$, by Lebesgue Diff. Thm.

we have $\int_{B(x,r)} |Du(z) - Du(x)|^p \rightarrow 0$ as $r \rightarrow 0$.

Consider $v(y) := u(y) - u(x) - Du(x) \cdot (y-x)$

$$\Rightarrow |u(y) - u(x) - Du(x) \cdot (y-x)| \lesssim r^{1-\frac{n}{p}} \left(\int_{B(x,2r)} |Du(x) - Du(z)|^p dz \right)^{\frac{1}{p}}$$

$$\lesssim r^{1-\frac{n}{p}} \cdot r^{\frac{n}{p}} \left(\int_{B(x,2r)} |Du(x) - Du(z)| dz \right)^{\frac{1}{p}}$$

$\rightarrow 0$ as $r \rightarrow 0^+$

Hence u is differentiable at x .

Now if $p = \infty$, note that $W_{loc}^{1,\infty} \subseteq W_{loc}^{1,p} \quad \forall \quad n < p < \infty$.

Corollary (Rademacher's Diff. Thm) u is locally Lipschitz continuous on U . Then u is differentiable a.e. on U .

RMK. The usual proof of Rademacher's Diff. Thm goes in two steps.

- Step 1 : 1 dimension case.
- 2 : n dimension

Hardy's inequality

Thm. ($n \geq 3$) Suppose $u \in H^1(B(0,r)) := W^{1,2}(B(0,r))$. Then $\frac{u}{|x|} \in L^2(B(0,r))$

$$\int_{B(0,r)} \frac{u^2}{|x|^2} dx \lesssim \int_{B(0,r)} |Du|^2 + \frac{u^2}{r^2} dx.$$

Proof. Assume $r=1$. $u \in C^\infty$.

Note $D(\frac{1}{|x|}) = -\frac{x}{|x|^3}$.

$$\int_{B(0,1)} \frac{u^2}{|x|^2} dx = - \int_{B(0,1)} u^2 D\left(\frac{1}{|x|}\right) \cdot \frac{x}{|x|} dx$$

$$= \int_{B(0,1)} \left[2u Du \cdot \frac{x}{|x|^2} + (n-1) \frac{u^2}{|x|^2} \right] dx - \int_{\partial B(0,1)} u^2 \underbrace{2 \cdot \frac{x}{|x|}}_{|x|=1 \text{ on } \partial B} dS$$

$$\Rightarrow (2-n) \int \frac{u^2}{|x|^2} dx = \underbrace{2 \int u Du \cdot \frac{x}{|x|^2} dx}_{\text{Cauchy Schwarz}} - \underbrace{\int_{\partial B(0,1)} u^2 dS}$$

$$= \int \operatorname{div}(xu^2) dx$$

$$= \int_{B(0,1)} nu^2 + 2u Du \cdot x$$

Thm. Characterization of H^s by Fourier transf.

Suppose $k \in \mathbb{N}$. (i) A function $u \in L^2(\mathbb{R}^n)$ belongs to $H^k(\mathbb{R}^n)$ iff $(1+|\xi|^2)^{k/2} \hat{u}(\xi) \in L^2(\mathbb{R}^n)$

(ii) For each $u \in H^k(\mathbb{R}^n)$, $\frac{1}{C_{k,n}} \|u\|_{H^k(\mathbb{R}^n)} \leq \|(1+|\xi|^2)^{k/2} \hat{u}(\xi)\|_{L^2} \leq C_{k,n} \|u\|_{H^k(\mathbb{R}^n)}$

Proof. 1. Assume $u \in H^k(\mathbb{R}^n) \Rightarrow D^\alpha u \in L^2(\mathbb{R}^n) \quad \forall |\alpha| \leq k$. Convention of Fourier Transf.

Suppose $u \in C_c^k(\mathbb{R}^n)$. Then we have $\widehat{D^\alpha u}(\xi) = (i\xi)^\alpha \hat{u}(\xi)$

$$\Rightarrow |\xi|^\alpha \hat{u}(\xi) \in L^2 \quad \forall |\alpha| \leq k.$$

$$\hat{f}(\xi) = \int_{\mathbb{R}^n} e^{-i\xi \cdot x} f(x) dx$$

$$f(x) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \hat{f}(\xi) e^{i\xi \cdot x} d\xi.$$

Choose $\alpha = (k, 0, \dots, 0), (0, k, 0, \dots, 0), \dots, (0, \dots, 0, k)$.

$$\Rightarrow \int_{\mathbb{R}^n} (1+|\xi|^2)^k \hat{u}(\xi) d\xi \leq C \|u\|_{H^k}.$$

By density, we could remove the assumption of $u \in C_c^k(\mathbb{R}^n)$

2. Conversely, if $(1+|\xi|^2)^{k/2} \hat{u} \in L^2(\mathbb{R}^n)$:

$$\| \underbrace{(i\xi)^\alpha \hat{u}}_{= \widehat{D^\alpha u}} \|_{L^2}^2 \leq \int_{\mathbb{R}^n} |\xi|^{2|\alpha|} |\hat{u}|^2 d\xi \lesssim \| (1+|\xi|^2)^{k/2} \hat{u} \|_{L^2}^2$$

$$\Rightarrow \widehat{D^\alpha u} \in L^2 \stackrel{\text{Plancherel}}{\Leftrightarrow} D^\alpha u \in L^2 \Rightarrow u \in H^k(\mathbb{R}^n)$$

□

Definition. Assume $s \in \mathbb{R}$. $u \in L^2(\mathbb{R}^n)$ We define

$$\|u\|_{H^s(\mathbb{R}^n)} := \| (1+|\xi|^2)^{s/2} \hat{u}(\xi) \|_{L^2(\mathbb{R}^n)} \quad \text{if it's finite.}$$

The Space of H^{-1} as the dual of $H_0^1(U)$.

Def. We denote $H^{-1}(U)$ the dual space of $H_0^1(U) := W_0^{1,2}(U)$

i.e. $f \in H^{-1}(U)$ if f is a bounded linear functional on $H_0^1(U)$.

RMK. We do not identify $H_0^1(U)$ with its dual,

Indeed, $H_0^1(U) \subset L^2 \subset H^{-1}(U)$

Notation: $\langle f, g \rangle$ pairing of $H^1(U)$ and $H_0^1(U)$

Def. For $f \in H^1(U)$, we define the norm

$$\|f\|_{H^1(U)} := \sup \left\{ \langle f, g \rangle \mid g \in H_0^1(U), \|g\|_{H_0^1(U)} \leq 1 \right\}$$

Theorem (Characterization of H^1) Assume $f \in H^1(U)$.

(i) $\exists$ functions $f^0, f^1, \dots, f^n$ in $L^2(U)$ s.t.

$$\langle f, v \rangle = \int_U f^0 v + \sum_{i=1}^n f^i \partial_{x_i} v \, dx \quad (*) \quad \forall v \in H_0^1(U)$$

$$(ii) \|f\|_{H^1(U)} = \inf \left\{ \left(\int_U \sum_{i=0}^n |f^i|^2 \, dx \right)^{\frac{1}{2}} \right\}$$

$$(iii) \langle v, u \rangle_{L^2(U)} = \langle v, u \rangle_{H^1} \quad \forall u \in H_0^1(U), v \in L^2(U) \subset H^1(U)$$

RMK. When $(*)$ holds, we often write $f = f^0 - \sum_{i=1}^n \partial_{x_i} f^i = \underbrace{L^2 + \text{div}(L^2\text{-vector function})}_{\text{function structure.}}$

Proof. (key idea: Riesz representation)

Given $u, v \in H_0^1(U)$, define $(u, v) := \int_U (Du \cdot Dv + uv) \, dx$, the inner product on $H_0^1(U)$

Let $f \in H^1(U)$. By Riesz, we can find a unique representative $u \in H_0^1(U)$ s.t. $\langle f, v \rangle = (u, v) \quad \forall v \in H_0^1(U)$

$$\int_U Du \cdot Dv + uv \, dx = \langle f, v \rangle \quad \forall v \in H_0^1(U)$$

$$\Rightarrow \begin{cases} f^0 = u \\ f^i = \partial_{x_i} u \end{cases} \quad (*) \text{ is then proved.}$$

let $u = v$.

$$(ii). \text{ Assume } f \in H^1(U) \quad \langle f, v \rangle = \int_U g^0 v + \sum g^i \partial_{x_i} v \, dx \quad \text{where } g^0, g^1, \dots, g^n \in L^2(U) \quad \text{is another characterization of } f.$$
$$\int_U |Du|^2 + u^2 \, dx = (u, u) \leq \int_U \sum_{i=0}^n |g^i|^2 \, dx$$

$$\Rightarrow \int_U \sum |f^i|^2 \, dx \leq \int_U \sum |g^i|^2 \, dx \quad \text{since } f^0 = u, f^i = \partial_{x_i} u.$$

The characterization from Riesz is the smallest one

$$3. |\langle f, v \rangle| \leq \left(\int_U \sum |f^i|^2 dx \right)^{\frac{1}{2}} \|v\|_{H_0^1(U)}$$

$$\Rightarrow \|f\|_{H_0^1(U)} \leq \left(\int_U \sum |f^i|^2 dx \right)^{\frac{1}{2}}$$

The equality holds if $v = \frac{u}{\|u\|_{H_0^1(U)}}$ $\leftarrow$ u is the representative of f in $H_0^1(U)$ by Riesz.



Elliptic Eqs.

$$\begin{cases} Lu = f & \text{in } U. \\ u = 0 & \text{on } \partial U \end{cases}$$

U is open bounded. f is given

$$Lu = - \sum_{i,j} a^{ij}(x) u_{x_i x_j} + \sum b^i(x) u_{x_i} + c(x) u \quad (\text{Divergence form})$$

OR

$$Lu = - \sum a^{ij}(x) u_{x_i} u_{x_j} + \sum b^i(x) u_{x_i} + c(x) u. \quad (\text{Non-div. form})$$

Assume the symmetry condition: $a^{ij} = a^{ji}$.

Def. We say L is (uniformly) Elliptic if $\exists \theta > 0$ s.t. $\sum a^{ij}(x) \xi_i \xi_j \geq \theta |\xi|^2 \quad \forall \xi \in \mathbb{R}^n, \text{ a.e. } x \in U$.

In particular, $(a^{ij}(x))_{i,j}$ is positive-definite, eigenvalues $\lambda_i \geq \theta$